

## **COVID-19 Immunization Schedule**

July 29, 2022

Booking a COVID-19 Vaccine Appointment, learn more.

Recommended Intervals for General Population and First Nation, Inuit, Metis								
Age Range/ Population Category	First Dose	Second Dose	Third Dose	Fourth Dose	Fifth Dose			
6 months – under 5	Can be received at any time	8 weeks after 1 <sup>st</sup> dose*	Not eligible	Not eligible	Not eligible			
5-11	Can be received at any time	8 weeks after 1 <sup>st</sup> dose*	Not eligible	Not eligible	Not eligible			
12-17	Can be received at any time	8 weeks after 1 <sup>st</sup> dose*	6 months (168 days) after 2 <sup>nd</sup> dose**	Not eligible	Not eligible			
18-59	Can be received at any time	8 weeks after 1 <sup>st</sup> dose*	5 months (140 days) after 2 <sup>nd</sup> dose***	5 months (140 days) after 3 <sup>rd</sup> dose***	Not eligible			
60+	Can be received at any time	8 weeks after 1 <sup>st</sup> dose*	5 months (140 days) after 2 <sup>nd</sup> dose***	5 months (140 days) after 3 <sup>rd</sup> dose***	Not eligible			
First Nation, Inuit, Metis Adults (18+) and their non- Indigenous household members	Can be received at any time	8 weeks after 1 <sup>st</sup> dose*	5 months (140 days) after 2 <sup>nd</sup> dose***	5 months (140 days) after 3 <sup>rd</sup> dose***	Not eligible			



Recommended Intervals for Immunocompromised Population									
Age Range/ Population Category	First Dose	Second Dose	Third Dose	Fourth Dose	Fifth Dose				
Immunocompromised due to health condition or medication AND First Nation, Inuit, Metis Adults (18+)	Can be received at any time	8 weeks after 1 <sup>st</sup> dose*	2 months (56 days) after 2 <sup>nd</sup> dose****	5 months (140 days) after 3 <sup>rd</sup> dose***	5 months (140 days) after 4 <sup>th</sup> dose***				
Immunocompromised due to health condition or medication (ages 6 months – under 5)	Can be received at any time	8 weeks after 1 <sup>st</sup> dose*	2 months (56 days) after 2 <sup>nd</sup> dose****	Not eligible	Not eligible				
Immunocompromised due to health condition or medication (ages 5-11)	Can be received at any time	8 weeks after 1 <sup>st</sup> dose*	2 months (56 days) after 2 <sup>nd</sup> dose****	Not eligible	Not eligible				
Immunocompromised due to health condition or medication (ages 12- 17)	Can be received at any time	8 weeks after 1 <sup>st</sup> dose*	2 months (56 days) after 2 <sup>nd</sup> dose****	6 months (168 days) after 3 <sup>rd</sup> dose**	6 months (168 days) after 4 <sup>th</sup> dose				
Immunocompromised due to health condition or medication (ages 18+)	Can be received at any time	8 weeks after 1 <sup>st</sup> dose*	2 months (56 days) after 2 <sup>nd</sup> dose****	5 months (140 days) after 3 <sup>rd</sup> dose***	5 months (140 days) after 4 <sup>th</sup> dose***				

\*Recommended time between doses. In discussion with your health care provider and/or vaccinator you may choose to receive this dose as early as 19 days for Pfizer, 21 days for Moderna, or 28 days for paediatric (6 months to 5 years) Moderna.

\*\* Recommended interval is 6 months (168 days) however may be received at the minimum interval of 3 months (84 days) after previous dose.

\*\*\*Recommended interval is 5 months (140 days) however may be received at the minimum interval of 3 months (84 days) after previous dose.

\*\*\*\* Recommended interval is 2 months (56 days) however may be received at the minimum interval of 28 days after previous dose.

**NOTE:** This chart is intended to provide high level guidance. Additional details on eligibility criteria can be found <u>here.</u>